

Technical Disclosure Commons

Defensive Publications Series

November 2021

Process For The Preparation Of Methyl (4,6-Diamino-2-(1-(2-Fluorobenzyl)-1H- Pyrazolo[3,4-B]Pyridin-3-Yl)Pyrimidin-5-Yl)Carbamate

MSN Laboratories Private Limited, R&D Center; Srinivasan Thirumalai Rajan; M Venkatesh;
Ramakrishna Bommidu; Balakrishna Gunupuru

Follow this and additional works at: https://www.tdcommons.org/dpubs_series

Recommended Citation

MSN Laboratories Private Limited, R&D Center; Srinivasan Thirumalai Rajan; M Venkatesh; Ramakrishna Bommidu; Balakrishna Gunupuru, "Process For The Preparation Of Methyl (4,6-Diamino-2-(1-(2-Fluorobenzyl)-1H-Pyrazolo[3,4-B]Pyridin-3-Yl)Pyrimidin-5-Yl)Carbamate", Technical Disclosure Commons, (November 15, 2021)
https://www.tdcommons.org/dpubs_series/4711




This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

This Article is brought to you for free and open access by Technical Disclosure Commons. It has been accepted for inclusion in Defensive Publications Series by an authorized administrator of Technical Disclosure Commons.

Process for the preparation of methyl (4,6-diamino-2-(1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b] pyridin-3-yl) pyrimidin-5-yl)carbamate

The present invention provides process for the preparation of methyl (4,6-diamino-2-(1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b] pyridin-3-yl)pyrimidin-5-yl)carbamate compound of formula-1, represented by the following structural formula:



Formula-1

Methyl(4,6-diamino-2-(1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl)pyrimidin-5-yl)carbamate is a process intermediate for the preparation of Riociguat.


Riociguat is a stimulator of soluble guanylate cyclase (sGC). It was approved by the U.S. Food and Drug Administration in October, 2013. It is used to treat two forms of pulmonary hypertension (PH): chronic thromboembolic pulmonary hypertension (CTEPH) and pulmonary arterial hypertension (PAH). Riociguat constitutes the first drug of the class of sGC stimulators.

US Patent No. 7173037 B2 discloses Methyl 4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-5-pyrimidinyl(methyl)carbamate and process for its preparation.

WO 03/095451 describes the preparation of Methyl (4,6-diamino-2-(1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl)pyrimidin-5-yl)carbamate compound of formula-1.

The present invention provides improved process for the preparation of Methyl (4,6-diamino-2-(1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl)pyrimidin-5-yl)carbamate of compound of formula-1, which provides Riociguat with high yield and purity. Also, the said process is cost effective and commercially viable.

Process of the present invention is schematically represented as below:



Examples:

Example-1: Preparation of methyl (4,6-diamino-2-(1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl)pyrimidin-5-yl)carbamate

Methanol [250 ml] and Tetrahydrofuran [250 ml] were added to (E)-2-(1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl)-5-(phenyldiazenyl)pyrimidine-4,6-diamine [100 g] at 25-30°C. Cooled the mixture to 5-10°C. Ammonium acetate [70.14 g] was added to the mixture at 5-10°C and stirred the mixture for 15 minutes. Zinc dust [36.35 g] was slowly added to the mixture at 5-10°C and stirred the mixture for 2 hours. Mixture filtered using hy-flow bed and bed washed with methanol. Slowly water (1000 ml) was added to the obtained filtrate at 25-30°C and stirred for 60 minutes. Adjusted the mixture pH between 2-3 by using hydrochloric acid at 25-30°C and stirred for 60 minutes. Filtered the solid and washed with water. Water (1000 ml) was added to the obtained solid at 25-30°C and adjusted the mixture pH between 8-9 by using sodium carbonate solution. Stirred the mixture for 60 minutes at 25-30°C. Filtered the solid and washed with water. Dried the compound at 60-65°C for 14 hrs.

Above obtained solid was added to the mixture of tetrahydrofuran (490 ml) and sodium bicarbonate solution (sodium bicarbonate 20.1 g and water 56 ml) at 25-30°C and stirred for 15 minutes. Cooled the mixture to 5-10°C. Slowly methyl chloroformate (22.6 g) was added to the mixture at 5-10°C and stirred for 5 hrs. Slowly water (1260 ml) was added to the mixture at 10-15°C. Raised the mixture temperature to 25-30°C and stirred for 16 hrs. Filtered the

precipitated solid and washed with water. Dichloromethane (1050 ml) and methanol (1050 ml) were added to the obtained wet solid at 25-30°C and stirred for 10 minutes. Under stirring carbon powder (28 g) and silica gel (14 g) were added to the mixture at 25-30°C and stirred for 2 hrs. Mixture filtered through hy-flow bed and bed washed with 1:1 solvent mixture of dichloromethane and methanol. Distilled-out the solvent from filtrate. Dimethylformamide (140 ml) was added to the mixture at 25-30°C. Heated the mixture to 70-75°C. Ethyl acetate (350 ml) was slowly added to the mixture at 70-75°C and stirred for 30 minutes. Cooled the mixture to 25-30°C and stirred for 2 hrs. Filtered the solid and solid washed with ethyl acetate (140 ml). Water (350 ml) was added to the obtained solid at 25-30°C. Heated the mixture to 45-50°C and stirred for 2 hours. Cooled the mixture to 25-30°C and stirred for 2 hrs. Filtered the solid and washed with water. Dried the obtained solid to get titled product.

Yield: 41.5 g; **Purity by HPLC:** 99.33%

Example-2: Preparation of 2-(1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl)pyrimidine-4,5,6-triamine

Methanol [250 ml] and Tetrahydrofuran [250 ml] were added to (E)-2-(1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl)-5-(phenyldiazenyl)pyrimidine-4,6-diamine [100 g] at 25-30°C. Cooled the mixture to 5-10°C. Ammonium acetate [70.14 g] was added to the mixture at 5-10°C and stirred the mixture for 15 minutes. Zinc dust [36.35 g] was slowly added to the mixture at 5-10°C and stirred the mixture for 2 hours. Mixture filtered using hy-flow bed and bed washed with methanol. Slowly water (1000 ml) was added to the obtained filtrate at 25-30°C and stirred for 60 minutes. Adjusted the mixture pH between 2-3 by using hydrochloric acid at 25-30°C and stirred for 60 minutes. Filtered the solid and washed with water. Water (1000 ml) was added to the obtained solid at 25-30°C and adjusted the mixture pH between 8-9 by using sodium carbonate solution. Stirred the mixture for 60 minutes at 25-30°C. Filtered the solid and washed with water. Dried the compound at 60-65°C for 14 hrs. to obtain titled product.

Yield; 71 g

Example-3: Preparation of methyl (4,6-diamino-2-(1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl)pyrimidin-5-yl)carbamate

2-(1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl)pyrimidine-4,5,6-triamine (71 g) was added to the mixture of tetrahydrofuran (490 ml) and sodium bicarbonate solution (sodium bicarbonate 20.1 g and water 56 ml) at 25-30°C and stirred for 15 minutes. Cooled the mixture to 5-10°C. Slowly methyl chloroformate (22.6 g) was added to the mixture at 5-10°C

and stirred for 5 hrs. Slowly water (1260 ml) was added to the mixture at 10-15°C. Raised the mixture temperature to 25-30°C and stirred for 16 hrs. Filtered the precipitated solid and washed with water. Dichloromethane (1050 ml) and methanol (1050 ml) were added to the obtained wet solid at 25-30°C and stirred for 10 minutes. Under stirring carbon powder (28 g) and silica gel (14 g) were added to the mixture at 25-30°C and stirred for 2 hrs. Mixture filtered through hy-flow bed and bed washed with 1:1 solvent mixture of dichloromethane and methanol. Distilled-out the solvent from filtrate. Dimethylformamide (140 ml) was added to the mixture at 25-30°C. Heated the mixture to 70-75°C. Ethyl acetate (350 ml) was slowly added to the mixture at 70-75°C and stirred for 30 minutes. Cooled the mixture to 25-30°C and stirred for 2 hrs. Filtered the solid and solid washed with ethyl acetate (140 ml). Water (350 ml) was added to the obtained solid at 25-30°C. Heated the mixture to 45-50°C and stirred for 2 hours. Cooled the mixture to 25-30°C and stirred for 2 hrs. Filtered the solid and washed with water. Dried the obtained solid to get titled product.

Yield: 41.5 g; **Purity by HPLC:** 99.33%
